

KAMIN V.S.

✓ Diagnostic value of the tellurite test in diphtheria. V.S.
Kamin (I. V. Stalin 2nd Moscow Med Inst). *Prilozhenie*
1954 1955. No. 5. 41-42. The tellurite test (Kamin's test)
is a rapid method for the detection of diphtheria. The
test is based on the fact that the tellurite test is
showing a positive result in about 24-48 hours and a
negative result in about 72-96 hours. The tellurite test
is a rapid method for the detection of diphtheria.
G. M. Kozlov

Clinic of Children's Infectious Diseases

KAZARIN V.S.
EXCERPTA MEDICA Sec.12 Vol.12/4 Ophthalmology April 58

647. CLINICAL FEATURES AND TREATMENT OF OCULAR DIPHTHERIA IN CHILDREN (Russian text) - Kazarin V.S. Clin. of Children's Infect. Dis. of the Sec. Med. Inst., Moscow - VOP. PEDIAT. 1956, 1/4 (7-13)

Forty-two patients with diphtheria of the eye were treated; most were received into the hospital on the 6th day and later on account of late diagnosis. The affection was unilateral in 26 patients, more often in the right eye. The catarrhal form of ocular diphtheria was seen in 4 patients, the croupous form in 36 and the dystrophic form in 2. The catarrhal form pursues the mildest clinical course, but is most

647

difficult to diagnose. The leading symptoms are: acute onset, rise in temperature, well-marked signs of toxæmia, palpebral oedema and redness, hyperæmia, swelling and tenderness of the conjunctiva, some difficulty in everting the lid, absence of photophobia, moderate sanguino-purulent discharge from the eye, rapid development of fairly thick glistening greyish-white films of variable size in the palpebral conjunctiva and injection of the scleral vessels. Localized or diffuse cloudiness of the cornea was observed rarely in the croupous form, but more or less constantly in the dystrophic form of diphtheria. In the treatment of the catarrhal form of ocular diphtheria the therapy may be confined to a single injection of 10,000-15,000 A. U. of antitoxic serum. In the case of the croupous and especially the diphtheritic forms, vigorous treatment is needed, viz.: repeated injections of serum (up to 40,000-80,000 A. U.), exhibition of tonics, penicillin injections and local therapy. In view of the fact that in some cases diphtheria bacilli were discovered in swabs not only from the eye but also from the pharynx and nose, repeated checking-up of swabs taken from the pharynx and nose is always necessary.

Lubenskaya - Leningrad (S)

KAZARIN, Vladimir Sergeyevich

[How to prevent angina in children] Kak predupredit' anginu
u detei. Moskva, Medgiz, 1958. 17 p.

(MIRA 13:12)

(RESPIRATORY ORGANS--DISEASES)

KAZARDI, V.S., kand. med. nauk

One of the methods for preventing biting through of the thread
in intubation patients. *Pediatrics* 36 no.11:77 N '58. (MIRA 12:8)

1. Iz kliniki detskikh infektsionnykh bolezney II Moskovskogo
meditsinskogo instituta imeni N.I. Pirogova na baze Detskoy
klinicheskoy bol'nitsy No.1.
(LARYNX--INTUBATION)

NISEVICH, N.I.; KAZARIN, V.S.

New advances in the treatment of anginas in children. Vop.okh.
mat.i det. 5 no.4:28-32 JI-Ag '60. (MIRA 13:7)

1. Iz kliniki detskikh infektsionnykh bolezney (sav. - prof.
D.D. Lebedev) II Moakovskogo meditsinskogo instituta im. N.I.
Pirogova (dir. - dotsent M.G. Sirotkina).
(TONSILS--DISEASES)

KAZARIN, V.S.; VERBENKO, A.A.; PROKHOROVA, L.V.

Diphtheria of the genitalia in girls and women. Vop. okh. mat. 1
det. 6 no. 2:83-85 F '61. (MIRA 14:2)

1. Iz kliniki detskikh infektsionnykh bolezney (zav. - zasluzhenny
deyatel' nauki prof. D.D. Lebedev) II Moskovskogo meditsinskogo
instituta imeni N.I. Pirogova (dir. - dotsent M.G. Sirotkina) i
polikliniki No. 3 Ministerstva zdravookhraneniya SSSR.

(GENERATIVE ORGANS, FEMALE—DISEASES) (DIPHTHERIA)

KAZARIN, V.S.; GASPARYAN, M.O.

Anginas in children caused by a yeastlike fungus from the genus
Candida. Sov. med. 25 no.2:38-42 F '62. (MIRA 15:3)

1. Iz kliniki detskikh infektsionnykh bolezney (zav. kafedroy -
prof. D.D. Lebedev) II Moskovskogo meditsinskogo instituta imeni
N.I. Pirogova (dir. - dotsent M.G. Sirotkina) na baze Detskoy
klinicheskoy bol'nitsy No.1 (glavnyy vrach Ye.M. Prokhorovich).
(MONILLIASIS)
(PHARYNX--DISEASES)

KAZARIN, V.S.

All-Union Conference on Hospital Service in the U.S.S.R. Vop.
okhr. materin. det. 8 no.1:91-93 '63 (MIRA 17:2)

GAABE, Yu.E.; KAZARINA, A.K.; KIPERMAN, G.Ya.; MALYI, I.G.;
ROZENTAL', O.E.; KOBOTKOV, A.F., retsenzent;
TITEL'BAUM, N.P., retsenzent; TRUKHANOVA, A.N., red.;
IL'YUSHENKOVA, T.P., tekhn. red.

[The theory of statistics] Teoriia statistiki. [By] IU.E.
Gaabe i dr. Pod red. I.G.Malogo. Moskva, Iskusstvo, 1963.
398 p. (MIRA 16:5)

(Statistics)

KAZARINA, A.N.

KAZARINA, A.N.; ANTROPOVA, V.N.

Effect of certain cortical and subcortical stimulants upon the development of streptococcal skin infections. Zhur.mikrobiol.epid.i immun. no.4:80 Ap '54. (MLRA 7:5)

1. Iz Odessakogo kozhno-venerologicheskogo instituta im. Glavche. (Narcotics) (Cerebral cortex) (Streptococcus)

KAZARINA, A.N.

USSR / Pharmacology, Toxicology. General Problems

U-1

Abs Jour : Ref. Zh.-Biol., No 2, 1958, No 7880

Author : Kazarina, A.N.

Inst :

Title : The Effect of Certain Pharmacological Agents Acting on the Central Nervous System on Streptococcal Skin Involvement in Rabbits.

Orig Pub : V. Sb.: Sovrem. vopr. dermatol., Kiyev, Gosmedizdat USSR, 1957, 23-27

Abstract : Experiments were performed on 90 rabbits. 1.2 billions microorganisms, obtained from a 24 hour culture of the hemolytic Streptococcus, were injected intracutaneously in the ears and thighs of all the animals. 75 rabbits were given various pharmacological agents daily during 1 month, while the

Card : 1/2

USSR/Microbiology - Antibiosis and Symbiosis. Antibiotics.

F-2

Abs Jour : Ref Zhur - Biologiya, No 7, 1957, 26304

Author : Belikov, G.P., Kudryavtseva, T.T., Antonova, A.A., Gugnayev, I.E., Kazarina, E.N.

Inst :

Title : Resistance of Dysentery Bacillus to Syntomycin, Streptomycin, and Biomycin (An Attempt at Comparative Study of Dyenteric Strains Isolated in 1953 in Moscow and Kishinev).

Orig Pub : Zh. mikrobiol., epidemiol., i immunobiologii, 1956, No 2, 35-41

Abst : Of the 800 strains of dysentery bacillus isolated in dysentery patients, 15.3% were found to be resistant to syntomycin (I). Most of the resistant strains were obtained from patients treated with I. Strains resistant to biomycin (II) and streptomycin (III) were not found. A comparative study of the sensitivity of

Card 1/2

Card 2/2

SHAYEVICH, A.B.; DANILEVSKAYA, V.V.; ZHOBOVA, N.I.; KAZARINA, G.P.;
TOROVINA, A.G.

Spectrographic determination of hydrogen in nickel and copper
and of oxygen in copper. Zav. lab. 30 no.11:1343-1346 '64
(MIRA 18:1)

1. Ural'skiy nauchno-issledovatel'skiy institut Chernykh
metallov.

KAZARINA, L. G.

"Experience in the Use of Tissue Therapy in Diseases of the Ear." Cand
Med Sci, Second Moscow State Medical Inst imeni I. V. Stalin, 1 Nov 54. (VM,
20 Oct 54)

Survey of Scientific and Technical Dissertations Defended at USSR Higher
Educational Institutions (10)

SO: Sum. 481, 5 May 55

CHAGIN, P.; TAVADZE, Ye.; FOMICHEV, N.; KAZARINA, N.

Material incentives and the quality of production; discussing the practice of the Kalinin textile workers. Sots.trud 7 no.7:113-121
Jl '62. (MIRA 15:8)

1. Nachal'nik otдела truda Pavlovo-Pokrovskoy fabriki (for Chagin). 2. Direktor Tbilisskogo kamvol'no-sukonnogo kombinata "Sovetskaya Gruzziya" (for Tavadze). 3. Direktor Shchelkovskogo khlopatobumazhnogo kombinata (for Fomichev). 4. Nachal'nik otдела truda Shchelkovskogo khlopatobumazhnogo kombinata (for Kazarina).

(Textile industry--Quality control)
(Bonus system)

LIVYY, G.V.; GAL'PEROVICH, M.G.; VASILYUK, N.Z.; SOPRIKO, A.Ye.;
KAZARINA, N.I.; CHURINA, V.I.; GIL'MAN, B.A.; YEGOROV, K.A.;
GONCHAR, Ye.G.

Method of refining the skin side of fur articles made with low
grade peltry; Soviet Certificate of Inventions No.147290. Kozh.-
obuv.prom. 4 no.8:43 Ag '62. (MIRA 15:8)
(Fur industry—Technological innovations)

KAZARINA, H.N.

Growing number of V.V. Gaganova's followers. Tekst. prom. 19
no.9:51-52 S '59. (MIRA 12:12)

1. Nachal'nik otдела truda Shchelkovskogo khlopchatobumazhnogo
kombinata.

(Efficiency, Industrial) (Textile industry)

KAZARINA, N.N.

Work practices after the introduction of a shorter workday.

Tekst.prom. 20 no.7:13-15 J1 '60.

(MIRA 13:7)

1. Nachal'nik otдела труда Shchelkovskogo khlopchatobumazhnogo kombinata.

(Shchelkovo--Textile factories)

KAZARINA, N.N., inzh.; GIL'MAN, B.A., inzh.; SLYUNIN, V.S., inzh.

New method of degreasing greasy sheep pelts. Izv.vys.ucheb.zav.;
tekh.leg.prom. no.2:29-32 '61. (MIRA 14:5)

1. Rekomendovana kafedroy tekhnologii kozhi Kiyevskogo tekhnologiche-
skogo instituta legkoy promyshlennosti.
(Hides and skins)

KAZARINA, N.N., inzh.; SHIFMAN, R.O., inzh.; GIL'MAN, B.A., inzh.;
RUDENKO, S.D., inzh.

Simplified method of determining the content of fatty substances
in leather and fur. Kozh.-obuv.prom. 4 no.8:28-29 Ag '62.
(MIRA 15:8)

(Leather) (Fur)

LIVYY, G.V.; KAZARINA, N.N.; GIL'MAN, B.A.; RUDENKO, S.D.; DREVINA, N.G.;
~~RESKERTNAYA~~, N.S.; ALPATSKAYA, V.P.; KOZLOVSKIY, S.I.;
SLYUNIN, B.S.

Development and application of reinforced film coating of sheepskins
for coats. Kozh.-obuv.prom. 4 no.3:25-28 Mr '62. (MIRA 15:5)
(Fur-Dressing and dyeing)

LIVYY, G.V., kand.tekhn.nauk; KAZARINA, N.N., inzh.; BRAGINSKIY, M.A., inzh.
SUKHOREBRYI, V.A., inzh.

Continuous diffusion of tanning materials in a rotating diffuser
unit. Nauch.-issl.trudy Ukr NIIKP no.13:68-76 '62.

(MIRA 18:2)

LIVYY, G.V., kand. tekhn. nauk; KAZARINA, N.N., inzh.; GIL'MAN, B.A., inzh.;
FASTOVETS, O.S., inzh.; MOROZYUK, N.I., inzh.; LITVINOV, Sh.I.,
inzh.; SAGAYDACHNYY, V.G., inzh.; BALAYFV, Ya.V., inzh.;
FITSA, A.S., inzh.

Manufacture of leather for lining and accessories from the
face split of DOL type pigskins. Kozh.-obuv. prom. 7 no.6:
29-32 Je '65. (MIRA 18:8)

KAZARINA, N.N.; SLUTSKIN, S.M.

Discussion on norms in the section of the Scientific and Technical
Society. Tekst. prom. 25 no.12:79-80 D '65. (MIRA 19:1)

LIVYY, G.V., kand. tekhn. nauk; FISH, B.I.; GORKUN, Yu.P.; LAZARINA, N.N.;
GIL'MAN, B.A.

Utilization of sheep pelts unsuitable for the production of fur
in the manufacture of chrome leather for shoe uppers and lining.
Kozh.-obuv. prom. 7 no.12:12-14 D '65.

(MIRA 19:2)

31547

188400

SOV/137-58-5-11661

Translation from: Referativnyy zhurnal, Metallurgiya, 1959, Nr 5, pp 311-312 (USSR)

AUTHORS: Rudnevskiy, N.K., Kozlova, N.V., Kazarina, T.P.

TITLE: Using a Spark and an Arc for Investigations Into the Dependence of the Intensity of Magnesium and Aluminum Lines on Their Concentration in a Binary Magnesium - Aluminum Alloy

PERIODICAL: Uch. zap. Gor'kovsk. un-ta, 1958, Nr 32, pp 161 - 167

ABSTRACT: The authors investigated the dependence of the absolute and the relative intensity of Mg and Al lines on their concentration in an Mg-Al alloy. A spark collected by the Rayskiy circuit and an a-c arc of the Sventitskiy circuit were used as sources for the spectrum excitation. Mg-Al alloys contained 2.3 - 9.8% Al. The specimens were bar-shaped having 1.5 x 4.5 x 3 cm dimension. The photographic records of the spectrum were made with an ISP-22 spectrograph. When analyzing Mg-Al alloys in the a-c arc, compared to the spark, changes in the current of its composition cause greater changes in the absolute intensity

Card 1/2

SOV/137-59-5-11661

Using a Spark and an Arc for Investigations Into the Dependence of the Intensity of Magnesium and Aluminum Lines on Their Concentration in a Binary Magnesium - Aluminum Alloy

of the alloy base lines, while the absolute intensity of Mg arc and spark lines remains practically unchanged in the spark of the investigated Mg concentration range, the changes in the arc are substantial. The character of changes in the intensity of Mg arc and spark lines is different. This may be explained by changes in the arc discharge temperature. It is shown that in the a-c arc the dependence of absolute and relative intensity of the Al I 3082.16 Å arc line on the Al concentration in the alloy (2 - 10%) is not described by Lomakin's formula, but by the exponential formula $J = Ae^{kc}$, where A and k are constant values, and c is the Al concentration in the alloy.

A.Sh.

Card 2/2

KAZARINOV, A.I. —

Example of the orthogenetic connection between gold ore formation and intrusions. *Biul. MOIP. Otd. geol.* 34 no.6:145-146
N-D '59. (MIRA 14:3)

(Aldan Plateau—Gold ores)

KAZARINOV, A. F.

5761. Sobol' Dal'nego Vostoka. Khabarovsk, Kn. izd., 1954. 120s. s ill. 20sm.
3.000 ekz 3r 35k. Bibliogr: s. 117-119-(55-1008) p. 639.113.5 (57.34) +(016.3)

SO: Knizhnaya, Ietopis, Vol. 1, 1955

KAZARINOV, A. P., Cond Biol Sci -- (diss) "Sable *Martes zibellina* L.
1758 of the Far East . (Systematics, ecology, biology, ^{Commercial} ~~trade~~ qualities,
^{restoration} ~~renewal~~ of ^{resources} ~~supplies~~, and their utilization)." Vladivostok, 1960.
16 pp (Far Eastern Affiliate in V.L. Komarov of the Siberian ^{Dept} ~~Branch~~
of the Acad of Sci USSR). 150 copies (EL,37,59, 107)

24

GURARI, Fabian Grigor'yevich; KAZARINOV, V.P., nauchnyy red.; KELLAREV, L.A.,
vedushchiy red.; GENNAD'YEV, I.M., tekhn. red.

[Geology, and oil and gas potentials of the Ob'-Irtysh interfluve]
Geologiya i perspektivy neftegazonosnosti Ob'-Irtyshskogo mezhdurech'ia.
Leningrad, Gos. nauchno.-tekhn. izd.-vo nef. i gornotoplivnoi lit-ry.
Leningr. otd-nie, 1959. 172 p. (Sibirskii nauchno-issledovatel'skii
institut geologii, geofiziki i mineral'nogo syr'ia. Trudy, no.3)

(Ob' Valley--Petroleum geology) (MIRA 12:12)

(Ob' Valley--Gas, Natural--Geology)

(Irtysh Valley--Petroleum geology)

(Irtysh Valley--Gas, Natural--Geology)

KAMENSHCHIKOV, Grigoriy Georgiyevich; KOLTUN, Sergey Ivanovich, inzh.;
NAUMOV, Vasilii Prokhorovich, inzh.; CHERNOBROVKIN, Boris
Sergeyevich, inzh.; POLYAKOV, V.P., inzh., retsenzent; KAZARINOV,
B.K., inzh., retsenzent; KON'KOV, A.S., dotsent, red.; DUGINA,
N.A., tekhn.red.

[Forging operations] Kuznechnoe proizvodstvo. Izd.3., 1apr. i
dop. Moskva, Gos.nauchno-tekhn.izd-vo mashinostroit.lit-ry, 1959.
447 p. (MIRA 12:8)

1. Uralmashzavod (for Koltun, Chernobrovkin). 2. Sverdlovskiy
zavod transportnogo mashinostroyeniya (for Naumov).
(Forging)

KAZARINOV, Boris Nikolayevich

KOLYUN. Sergey Ivanovich; KAZARINOV, Boris Nikolayevich; KAYDALOV, P.K.,
inzhener, spetsent; DUGINA, N.A., tekhnicheskiy redaktor.

[Improvements in forge shops; practices of the Ural Machine
Manufacturing Plant] Usovershenstvovaniia v kuznechnom tsekhe;
iz opyta Uralsmashsavyoda. Moskva, Gos.nauchno-tekhn.isd-vo
mashinostroit.lit-ry, 1956. 51 p. (MIRA 10:6)
(Forging machinery)

KAZARINOV, B.N., inzhener.

Blacksmith M.N. Kushev. Mashinostroitel' no.11:33-34 N '57.
(MIRA 10:10)

(Kuskov, Mikhail Nikhanorovich)

KAZARINOV B.N.

PHASE I BOOK EXPLOITATION 1042

Ural'skiy zavod tyazheloego mashinostroyeniya, Sverdlovsk

Kovka i termicheskaya obrabotka (Forging and Heat Treatment) Moscow, Mashgiz, 1958. 132 p. (Series: Its Sbornik statey, vyp 5) 6,000 copies printed.

Ed.: Kvater, I.S., Engineer; Tech. Ed.: Dugina, N.A.; Ed. (Ural-Siberian Division, Mashgiz): Sustavov' M.I., Engineer.

PURPOSE: This book is intended for engineers and technicians working in the field of forging and heat-treating of metals.

COVERAGE: The book presents material which reflects the achievements of Uralmashzavod (Ural Heavy Machine-building Plant imeni S. Ordzhonikidze) in the field of forging and heat-treating of metals. Various improvements in production methods, mechanization and automation of forging and heat-treating processes, application of various methods of inspection of forgings and elimination of rejects are described. Specific information on improvements in

Card 1/4

APPROVED FOR RELEASE: 06/13/2000
Forging and Heat Treatment 1042

CIA-RDP86-00513R000721320018-7"

forging and heat-treating of large parts such as turbine discs and rotors, cold-rolling-mill rolls, and crankshafts are presented. Descriptions are given of the results of new studies undertaken with a view to elimination of rejects and improvement of the quality of parts, determination of residual stresses at various cooling speeds, data on the efficiency of ultrasonic inspection and the effect of degassing of molten steel on the quality of forgings. The book was prepared by the members of the plant organization of NTOMashprom in connection with the 25th anniversary of the Ural Heavy Machine-building Plant.

TABLE OF CONTENTS:

Kvater, I.S. Summary of Development of Forging and Heat-treating Production at the Ural Heavy Machine-building Plant	3
Lebedev, A.V., and Ustyugov, P.A. Welding Up of Internal Flaws in Large Forgings	21
Zlatkin, M.G. Improvement of Open-die Forging	34

Card 2/4

Forging and Heat Treatment	1042	46
Katkov, I.S. Improvement of Methods of Forging Turbine-discs		50
Kazarinov, B.N. Comprehensive Mechanization of Press-forging Processes		59
ABSTRACT: Measures undertaken to mechanize difficult and labor-consuming processes are described together with the specifications, the design, and the performance of the following facilities: A floor-level forging manipulator; a mechanism permitting automatic control of the drop hammer; an electrical buggy equipped with a turntable; combined roll table supports for forging of rings; special cover plates for the gripping jaws of the manipulator to facilitate the gripping of billets during forging of strip.		73
Mikul'chik, A.V., and Kats, Sh. I. Schistose Type of Fracture in Chrome-nickel-molybdenum Steel		90
Card 3/4		103

- Forging and Heat Treatment 1042
- Zabludovskiy, V.M. Effect of the Speed of Cooling of Cylindrical
Parts on the Magnitude of Residual Stresses 111
- Zabludovskiy, V.M. Determination of Residual Stresses in
Large Parts 115
- Kozhevnikov, M.A. Investigation of Parts Rejected on
Ultrasonic Inspection 123
- Kuruklis, G.L., and Vereshchagina, M.G. Sulphidization of Machine
Parts 130

AVAILABLE: Library of Congress

GO/ksv
1-7-59

Card 4/4

AUTHOR: Kazarinov, B.N., Engineer SOV/122-58-6-14/37
TITLE: ~~Examples of Mechanisation~~ of Hand-forging Operations
at the Ural Machinery Plant (Primery mekhanizatsii svobodnoy
kovki na Uralskoy zavode)

PERIODICAL: Vestnik Mashinostroyeniya, 1958, Nr 6, pp 41-42 (USSR)

ABSTRACT: Examples are illustrated of manipulating and transporting equipment designed and built for the mechanisation of hand-forging operations. These include an electrically propelled carriage with a rotary table mounted on balls, a 2-ton manipulator used in the rolling of rings from forged and pierced blanks and a 2-ton gripping manipulator, controlled by a single operator. There are 4 figures.

Card 1/1

1. Industrial plants--Equipment 2. Materials--Handling

SHESTAKOV, Andrian Andrianovich; LEVANDOVSKIY, P.G., inzh., retsenzent;
KAZARINOV, B.W., inzh., red.; PUCHKOV, S.G., inzh., red.;
DUGINA, N.A., tekhn.red.

[Steam and pneumatic hammer operator] Mashinist parovozdushnogo
molota. Moskva, Gos.nauchno-tekhn.izd-vo mashinostroit.lit-ry,
1959. 118 p. (MIRA 12:4)

(Forging machinery)

DATE 10/2/82

[illegible]

Matematika i arifmetika matematika i arifmetika
atom and arithmetic in the mathematics
1973. 519 p. 12,000 copies printed.

MR. B. V. FALLON, Doctor of Technical Sciences, Chief of the Metallurgical Department, P. P. Polzunov Scientific Institute of Technical Sciences, 26, V. O. District, Leningrad, U. S. S. R.

MR. A. A. KROKHIN, Doctor of Technical Sciences, Chief of the Metallurgical Department, P. P. Polzunov Scientific Institute of Technical Sciences, 26, V. O. District, Leningrad, U. S. S. R.

MR. A. I. KROKHIN, Candidate of Technical Sciences, Chief of the Metallurgical Department, P. P. Polzunov Scientific Institute of Technical Sciences, 26, V. O. District, Leningrad, U. S. S. R.

MR. A. I. KROKHIN, Candidate of Technical Sciences, Chief of the Metallurgical Department, P. P. Polzunov Scientific Institute of Technical Sciences, 26, V. O. District, Leningrad, U. S. S. R.

[illegible]

Continues and Discontinues Paying Into Bonds (Continued)

NAME	DATE	AMOUNT	REMARKS
Buttner, M. B. Lyons, Engineer, and A. D. Popov, Mechanical Engineer	1914	100.00	Continued
	1915	100.00	Continued
	1916	100.00	Continued
	1917	100.00	Continued
	1918	100.00	Continued
	1919	100.00	Continued
	1920	100.00	Continued
	1921	100.00	Continued
	1922	100.00	Continued
	1923	100.00	Continued
	1924	100.00	Continued
	1925	100.00	Continued
	1926	100.00	Continued
	1927	100.00	Continued
	1928	100.00	Continued
	1929	100.00	Continued
	1930	100.00	Continued
	1931	100.00	Continued
	1932	100.00	Continued
	1933	100.00	Continued
	1934	100.00	Continued
	1935	100.00	Continued
	1936	100.00	Continued
	1937	100.00	Continued
	1938	100.00	Continued
	1939	100.00	Continued
	1940	100.00	Continued
	1941	100.00	Continued
	1942	100.00	Continued
	1943	100.00	Continued
	1944	100.00	Continued
	1945	100.00	Continued
	1946	100.00	Continued
	1947	100.00	Continued
	1948	100.00	Continued
	1949	100.00	Continued
	1950	100.00	Continued
	1951	100.00	Continued
	1952	100.00	Continued
	1953	100.00	Continued
	1954	100.00	Continued
	1955	100.00	Continued
	1956	100.00	Continued
	1957	100.00	Continued
	1958	100.00	Continued
	1959	100.00	Continued
	1960	100.00	Continued
	1961	100.00	Continued
	1962	100.00	Continued
	1963	100.00	Continued
	1964	100.00	Continued
	1965	100.00	Continued
	1966	100.00	Continued
	1967	100.00	Continued
	1968	100.00	Continued
	1969	100.00	Continued
	1970	100.00	Continued
	1971	100.00	Continued
	1972	100.00	Continued
	1973	100.00	Continued
	1974	100.00	Continued
	1975	100.00	Continued
	1976	100.00	Continued
	1977	100.00	Continued
	1978	100.00	Continued
	1979	100.00	Continued
	1980	100.00	Continued
	1981	100.00	Continued
	1982	100.00	Continued
	1983	100.00	Continued
	1984	100.00	Continued
	1985	100.00	Continued
	1986	100.00	Continued
	1987	100.00	Continued
	1988	100.00	Continued
	1989	100.00	Continued
	1990	100.00	Continued
	1991	100.00	Continued
	1992	100.00	Continued
	1993	100.00	Continued
	1994	100.00	Continued
	1995	100.00	Continued
	1996	100.00	Continued
	1997	100.00	Continued
	1998	100.00	Continued
	1999	100.00	Continued
	2000	100.00	Continued
	2001	100.00	Continued
	2002	100.00	Continued
	2003	100.00	Continued
	2004	100.00	Continued
	2005	100.00	Continued
	2006	100.00	Continued
	2007	100.00	Continued
	2008	100.00	Continued

**2025 30 Year Bond Redemption
Surrender Notice
(Section 1)**

7. Mechanization of Production for Pure Profit-making Enterprises
Examples of mechanization of various operations
Automation of processing of casting sand
Mechanization of lines for sheet-rolling

8. Special Casting Methods (Chernozvov, N.Ye., and E.V. Knyalov, Engineers)

9. Pneumatic Transport of Loose Materials (Vertikalbau, No. 2, p. 287).
Friedrich and P. A. Schmitt
Lavalentz und Frickhagen

FOR CREDIT IN HISTORY OR CIVILIZATION

1. Full Notification of Possible Perings (Leverability, P.D., and D.I.
Insulation, Engineers)

Use of simulators in the making of foreign
currency equipment for the mechanization of
mining

System of Machines and Mechanisms for Stamping (Zachary, A.E., Ed.,
New York, V.P. Massey, E.J. Probst, and Paul Zurever,
Editors), 1967, 1968, 1969, 1970, 1971, 1972, 1973, 1974, 1975, 1976, 1977, 1978, 1979, 1980, 1981, 1982, 1983, 1984, 1985, 1986, 1987, 1988, 1989, 1990, 1991, 1992, 1993, 1994, 1995, 1996, 1997, 1998, 1999, 2000, 2001, 2002, 2003, 2004, 2005, 2006, 2007, 2008, 2009, 2010, 2011, 2012, 2013, 2014, 2015, 2016, 2017, 2018, 2019, 2020, 2021, 2022, 2023, 2024, 2025, 2026, 2027, 2028, 2029, 2030, 2031, 2032, 2033, 2034, 2035, 2036, 2037, 2038, 2039, 2040, 2041, 2042, 2043, 2044, 2045, 2046, 2047, 2048, 2049, 2050, 2051, 2052, 2053, 2054, 2055, 2056, 2057, 2058, 2059, 2060, 2061, 2062, 2063, 2064, 2065, 2066, 2067, 2068, 2069, 2070, 2071, 2072, 2073, 2074, 2075, 2076, 2077, 2078, 2079, 2080, 2081, 2082, 2083, 2084, 2085, 2086, 2087, 2088, 2089, 2090, 2091, 2092, 2093, 2094, 2095, 2096, 2097, 2098, 2099, 2100, 2101, 2102, 2103, 2104, 2105, 2106, 2107, 2108, 2109, 2110, 2111, 2112, 2113, 2114, 2115, 2116, 2117, 2118, 2119, 2120, 2121, 2122, 2123, 2124, 2125, 2126, 2127, 2128, 2129, 2130, 2131, 2132, 2133, 2134, 2135, 2136, 2137, 2138, 2139, 2140, 2141, 2142, 2143, 2144, 2145, 2146, 2147, 2148, 2149, 2150, 2151, 2152, 2153, 2154, 2155, 2156, 2157, 2158, 2159, 2160, 2161, 2162, 2163, 2164, 2165, 2166, 2167, 2168, 2169, 2170, 2171, 2172, 2173, 2174, 2175, 2176, 2177, 2178, 2179, 2180, 2181, 2182, 2183, 2184, 2185, 2186, 2187, 2188, 2189, 2190, 2191, 2192, 2193, 2194, 2195, 2196, 2197, 2198, 2199, 2200, 2201, 2202, 2203, 2204, 2205, 2206, 2207, 2208, 2209, 2210, 2211, 2212, 2213, 2214, 2215, 2216, 2217, 2218, 2219, 2220, 2221, 2222, 2223, 2224, 2225, 2226, 2227, 2228, 2229, 2230, 2231, 2232, 2233, 2234, 2235, 2236, 2237, 2238, 2239, 2240, 2241, 2242, 2243, 2244, 2245, 2246, 2247, 2248, 2249, 2250, 2251, 2252, 2253, 2254, 2255, 2256, 2257, 2258, 2259, 2260, 2261, 2262, 2263, 2264, 2265, 2266, 2267, 2268, 2269, 2270, 2271, 2272, 2273, 2274, 2275, 2276, 2277, 2278, 2279, 2280, 2281, 2282, 2283, 2284, 2285, 2286, 2287, 2288, 2289, 2290, 2291, 2292, 2293, 2294, 2295, 2296, 2297, 2298, 2299, 2300, 2301, 2302, 2303, 2304, 2305, 2306, 2307, 2308, 2309, 2310, 2311, 2312, 2313, 2314, 2315, 2316, 2317, 2318, 2319, 2320, 2321, 2322, 2323, 2324, 2325, 2326, 2327, 2328, 2329, 2330, 2331, 2332, 2333, 2334, 2335, 2336, 2337, 2338, 2339, 2340, 2341, 2342, 2343, 2344, 2345, 2346, 2347, 2348, 2349, 2350, 2351, 2352, 2353, 2354, 2355, 2356, 2357, 2358, 2359, 2360, 2361, 2362, 2363, 2364, 2365, 2366, 2367, 2368, 2369, 2370, 2371, 2372, 2373, 2374, 2375, 2376, 2377, 2378, 2379, 2380, 2381, 2382, 2383, 2384, 2385, 2386, 2387, 2388, 2389, 2390, 2391, 2392, 2393, 2394, 2395, 2396, 2397, 2398, 2399, 2400, 2401, 2402, 2403, 2404, 2405, 2406, 2407, 2408, 2409, 2410, 2411, 2412, 2413, 2414, 2415, 2416, 2417, 2418, 2419, 2420, 2421, 2422, 2423, 2424, 2425, 2426, 2427, 2428, 2429, 2430, 2431, 2432, 2433, 2434, 2435, 2436, 2437, 2438, 2439, 2440, 2441, 2442, 2443, 2444, 2445, 2446, 2447, 2448, 2449, 2450, 2451, 2452, 2453, 2454, 2455, 2456, 2457, 2458, 2459, 2460, 2461, 2462, 2463, 2464, 2465, 2466, 2467, 2468, 2469, 2470, 2471, 2472, 2473, 2474, 2475, 2476, 2477, 2478, 2479, 2480, 2481, 2482, 2483, 2484, 2485, 2486, 2487, 2488, 2489, 2490, 2491, 2492, 2493, 2494, 2495, 2496, 2497, 2498, 2499, 2500, 2501, 2502, 2503, 2504, 2505, 2506, 2507, 2508, 2509, 2510, 2511, 2512, 2513, 2514, 2515, 2516, 2517, 2518, 2519, 2520, 2521, 2522, 2523, 2524, 2525, 2526, 2527, 2528, 2529, 2530, 2531, 2532, 2533, 2534, 2535, 2536, 2537, 2538, 2539, 2540, 2541, 2542, 2543, 2544, 2545, 2546, 2547, 2548, 2549, 2550, 2551, 2552, 2553, 2554, 2555, 2556, 2557, 2558, 2559, 2560, 2561, 2562, 2563, 2564, 2565, 2566, 2567, 2568, 2569, 2570, 2571, 2572, 2573, 2574, 2575, 2576, 2577, 2578, 2579, 2580, 2581, 2582, 2583, 2584, 2585, 2586, 2587, 2588, 2589, 2590, 2591, 2592, 2593, 2594, 2595, 2596, 2597, 2598, 2599, 2600, 2601, 2602, 2603, 2604, 2605, 2606, 2607, 2608, 2609, 2610, 2611, 2612, 2613, 2614, 2615, 2616, 2617, 2618, 2619, 2620, 2621, 2622, 2623, 2624, 2625, 2626, 2627, 2628, 2629, 2630, 2631, 2632, 2633, 2634, 2635, 2636, 2637, 2638, 2639, 2640, 26

Standard arrangement for lifting, turning, and transferring of blocks	122
Mechanization and automation of auxiliary operations	122
Full mechanization of production lines	130
	131

mechanization in production lines
Mechanization and Automation of Die-Forging Equipment (Mikhlin, I.I.,
M.L. Alekseyev, L.M. Kabanov, B.L. Dobryshin, P.N. Ponomarev, and
A.A. Vorobeyev, Engineers)
127

161	Automation of the thermal processes in heating furnaces
162	Examples of modernization of equipment
163	Automatic and Semi-automatic Dies [with automatic feed and ejection] (Institute, S.V., Candidates of Technical Sciences and N.Y. Potekhin, Engineer)
164	

156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974

KOLTUN, Sergey Ivanovich; BORINSKIY, Mikhail L'vovich; KATKOV, Leonid Ivanovich; KAZARINOV, Boris Nikolayevich; KATKOV, N.P., inzh., retsenzent; BASSEYN, V.V., inzh., retsenzent; KATKOV, I.S., inzh., red.; YERMAKOV, N.P., tekhn.red.

[Mechanization of minor processes in press forging plants]
Malaia mekhanizatsiia kuznechno-pressovykh tsakhov; al'bom
chartezhei. Moskva, Gos.nauchno-tekhn.isd-vo mashinostroit.
lit-ry, 1960. 104 p. (MIRA 14:2)
(Forge shops--Equipment and supplies)

KAZARINOV, B.N.

"Mechanisms, devices and means of mechanizing press forging operations" by V.N. Smirnov, P.M. Pavlovich. Reviewed by B.N.Kazarinov.
Kuz.-shtam. proizv. 3 no.11:46-47 N '61. (MIRA 14:11)
(Forging machinery) (Sheet metal working machinery)
(Smirnov, V.N.) (Pavlovich, P.M.)

ACCESSION NR: AP4034598

S/0182/64/000/004/0005/0007

AUTHORS: Kazarinov, B. N.; Shaburov, V. Ye.

TITLE: Investigation of the process of closing axial defects by upsetting

SOURCE: Kuznechno-ditampovochnoye proizvodstvo, no. 4, 1964, 5-7

TOPIC TAGS: forging, defect structure, defect formation, lead, steel, steel mill/
UIM50 testing machine, U7 steel

ABSTRACT: The authors proposed and elaborated the method of upsetting for use in closing axial defects in steel and lead. The influence of nonhomogeneity of deformation on both the closing of defects and on the strain condition of the sample and the dependence of defect closing upon sample size and form were also investigated. The samples (made from U7 steel and white lead) were cylindrical, with a circular orifice cut into the axis of each sample. Testing was carried out with testing machine UIM-50; samples were placed in a special container (see Fig. 1 on the Enclosure) for use in conjunction with the testing machine. A photographic record shows the sequential steps in the closing of defects and gives recordings of the change in H/D ratio. The authors present a schematic diagram showing the mechanism of defect closing. It is concluded that: 1) the magnitude of the H/D

Card 1/3

ACCESSION NR: AP4034598

ratio is the basic factor influencing axial defect closure, 2) defect closure occurs irregularly, beginning in the defect center and distributing itself to the contact surface due to nonhomogeneous deformation, and 3) with increasing non-homogeneity of deformation, caused by contact friction, defect closure occurs more intensely and increases with greater H/D ratio. Orig. art. has: 3 figures, 4 photographs, and 1 equation.

ASSOCIATION: none

SUBMITTED: 00

SUB CODE: MT, MM

NO REF SOV: 003

ENCL: 01

OTHER: 001

Card 2/3

ACCESSION NR: AP4034598

ENCLOSURE: 01




Fig. 1.
Container

Card 3/3

L 20241-AR
DPT(M)/SWP(W)/FWA(d) / T(SWP) : ...
ACCESS : ...

Судья К. И. Тетерин г. Яд.

improved technology for forging E14078, and
Sht 6: Ruznechno-shtempovachnye proizvozhitel'stva

alloy, E143740 alloy, Nitrogen
alloy, the shape, forging technique, mechanical.

3.1.2. Harmonic 89A (type) alloy were cast in 100 g
3.1.3. The alloy was used in the experiments. The
3.1.4. The alloy was used in the experiments. The
3.1.5. The alloy was used in the experiments. The
3.1.6. The alloy was used in the experiments. The
3.1.7. The alloy was used in the experiments. The
3.1.8. The alloy was used in the experiments. The
3.1.9. The alloy was used in the experiments. The
3.1.10. The alloy was used in the experiments. The

L 2041-05

ACCESSION NP: AP5001777

In 20—25 hr and forged into 220-mm round or 200 x 2 3/4 mm
billets without reheating. The forging was done by
forged by the above technique had a tensile strength
of 100,000 psi, a reduction of area of 10%
and a yield strength of 50,000 psi. The material was
used for the manufacture of 220-mm rounds and 200 x 2 3/4 mm
billets. The material was used for the manufacture of 220-mm
rounds and 200 x 2 3/4 mm billets. The material was used for the
manufacture of 220-mm rounds and 200 x 2 3/4 mm billets.

ASSOCIATION: none

SUBMITTED: 00

ENCL: 00

SUB CODE: 44

NO REF SOV: 000

OTHER: 000

ATD PRESS: 121

Card 2/2

L 20776-66 EWT(m)/EWP(w)/EWA(d)/EWP(v)/T/EWP(t)/EWP(k)/ETC(m)-6 JD/HW/FM	
ACC NR: AP6004679	SOURCE CODE: UR/0182/65/000/010/0001/0006
AUTHOR: <u>Kazarinov, B. N.; Okhrimenko, Ya. M.</u>	
ORG: none	
TITLE: <u>Improvements in the process of forging turbine disks from EI437BU alloy</u>	
SOURCE: <u>Kuznechno-shtampovoychnoye proizvodstvo, no. 10, 1965, 1-6</u>	
TOPIC TAGS: turbine disk, drop forging, hot forging, metal grain structure / EI437BU Cr-Ni high-temperature alloy, solid mechanical property	
<p>ABSTRACT: The turbine disks forged from EI437BU Cr-Ni high-temperature alloy display some nonuniformity of mechanical properties and stress-rupture strength owing chiefly to their consertal structure; this may be offset by properly adjusting the forging technology, the paramount objective being to reduce to a minimum the danger of the formation of internal cracks in the finished elements (turbine disks). In this connection, the authors describe new experimental techniques for processing 700-kg ingots of EI437BU alloys into turbine disks. The blanks used are round rather than square, and this eliminates the possibility of the rise of axial cracks during the rounding process. Moreover, the hot upsetting of round blanks proceeds more uniformly, particularly when their height-to-diameter ratio is <2.5. Following their hot upsetting the blanks are heated to 1150+10°C in two-chamber gas furnaces and then drop-forged in dies heated to 250-300°C and lubricated with dry sawdust. The principle</p>	
Card 1/2	UDC: 621.73.032

L 20775-66

ACC NR: AP6004679

pal advantage of the new technique appears to be the replacement of square blanks with round blanks, since comparison tests showed that disks forged from round blanks (diameter 220 mm) display superior mechanical properties and more uniform macro-structure before and after heat treatment (hardening at 1080°C for 8 hr with cooling in air; aging at 750±5°C for 16 hr with cooling in air) compared with the disks forged from square blanks (200x200 mm), since ready-made round blanks are free of the stresses present in square blanks when these have to be rounded prior to their hot upsetting. Orig. art. has: 7 figures, 2 tables.

SUB CODE: 11, 13, 20/ SUB DATE: none/ ORIG REF: 006/ OTH REF: 000

Card

2/2

vmb

KAZARINOV, F.

New equipment for geologists. MTO no.4:21 Ap '59.

(MIRA 12:6)

1. Chlen prezidiuma Tsentral'nogo pravleniya nauchno-tekhnicheskogo
gornogo obshchestva, (Geological research)

...KAZARINOV, F.F.; SMIRNOV, A.I.

Problems of trade-union organizations of workers employed in geological prospecting as presented by the Third Trade-Union Congress.
Sov.geol. 1 no.9:3-12 S '58.
(MIRA 12:2)

1. TSentral'nyy Komitet profsoyuza rabochikh geologorazvedochnykh
rabot.

(PROSPECTING)

AUTHOR: Kazarinov, F.F.

SOV-132-50-9-18/10

TITLE: New Safety Rules for Geological Prospecting Operations (Novyye yedinye pravila bezopasnosti pri geologorazvedochnykh rabotakh)

PERIODICAL: Razvedka i okhrana nedr, 1958, ¹³Nr 9, pp 60-62 (USSR)

ABSTRACT: The author describes the new safety rules for geological and prospecting operations and stresses the importance of their strict observance by all concerned. New courses must be created for the inspectors of work safety so that these rules can be successfully applied.

ASSOCIATION: TsK Profsoyuza rabochikh geologorazvedochnykh rabot (Central Committee fo the Trade Union of Geological Workers)

1. Geophysical prospecting--Safety measures

Card 1/1

USCOMM-DC-55791

KAZARINOV, F.F.

Improve working conditions. Razved. i okh. nedr 26 no.2:
49-51 Feb. '60. (MIRA 14:6)

1. Tsentral'nyy komitet profsoyuza rabochikh geologorazvedochnykh
rabot.

(Prospecting--Safety measures)

KAZARINOV, F.F.

For all-out sanitation and improvement of working conditions.
Razved.i okh.nedr 28 no.4:51-54 Ap '62. (MIRA 15:4)

1. TSentral'nyy komitet profsoyuza rabochikh geologorazvedochnykh
rabot.

(Prospecting--Safety measures)

"APPROVED FOR RELEASE: 06/13/2000

CIA-RDP86-00513R000721320018-7

APPROVED FOR RELEASE: 06/13/2000

CIA-RDP86-00513R000721320018-7"

KAZANOV, I. A.

Seleniye vypritsmiteli dlia predpriatii svyazi /Selenium rectifiers for communications enterprises/. Moskva, Svyaz'izdat, 1952. 252 p.

SO: Monthly List of Russian Accessions, Vol 6 No 4, July 1953

KAZARINOV, Ivan Alekseyevich; KOKOSHKIN, Pavel Aleksandrovich; KULESHOV,
V.N., otv.red.; KONDRASHINA, N.M., red.; MARKOCH, K.G., tekhn.red.

[Design of power supply devices for wire-communication enterprises]
Proektirovaniye elektropitalushchikh ustanovok predpriyatii pro-
vodnoi svyazi. Moskva, Gos.izd-vo lit-ry po voprosam svyazi i
radio, 1960. 399 p. (MIRA 14:3)

(Electric power supply for apparatus)
(Telegraph) (Telephone)

BOVKUN, Viktor Georgiyevich; KAZARINOV, Ivan Alekseyevich; KOKOSHKIN, Pavel Aleksandrovich; LYUBSKIY, Gennadiy Severianovich; MEDOVAR, Anatoliy Isayevich; PETROV, Viktor Vasil'yevich; PIONTKOVSKIY, Bronislav Aleksandrovich; SERDYAKOV, Nikolay Ivanovich; ELINSON, Mikhail Mikhaylovich; SERGEYCHUK, K.Ya., red.; GRIGOR'YEV, B.S., red.; FORTUSHENKO, A.D., red.; BUSANKINA, N.G., red.; SHEFER, G.I., tekhn. red.

[Engineering manual on electric communications; electric equipment] Inzhenerno-tekhnicheskii spravochnik po elektrosviazi; elektroustanovki. Moskva, Gos. izd-vo lit-ry po voprosam sviazi i radio, 1962. 671 p. (MIRA 15:6)
(Telecommunication--Handbooks, manuals, etc.)
(Electric engineering--Handbooks, manuals, etc.)

KAZARINOV, J.

SCIENCE

PERIODICALS: ~~ACTA ZOOLOGICA~~. Vol. 3, no. 4, 1955
MAGYAR FIZIKAI FOLYOIRAT. Vol. 3, no. 4, 1955.

Kazarinov, J. Elastic scattering of protons on 380-Mev energy protons. Tr.
from the Russian. p. 427

Monthly list of East European Accessions (EEAI) LC, Vol. 8, No. 2
February 1959, Unclass.

HAZARENOV, K. A. (Eng.); SHITS, Yu. R. (Cand. Tech. Sci.)

"Questions of the Application of Telemechanics in Communications,"

paper read at the Session of the Acad. Sci. USSR, on Scientific Problems of Automatic
Production, 15-20 October 1956.
Avtomatika i telemekhanika, No. 2, p. 182-192, 1957.

9015229

GURVICH, S.I.; KAZARINOV, L.N.; MALASHEVSKIY, A.N.

Discovery of titanium-zirconium placers in central Ciscaucasia.

Dokl.AN SSSR 144 no.3:619-621 My '62.

(MIRA 15:5)

1. Predstavleno akademikom D.I.Shcherbakovym.
(Ciscaucasia—Geology, Stratigraphic)

GURVICH, S.I.; KAZARINOV, I.N.; KHMARA, N.V.

[Ancient rare-metal-titanium placers, methods of prospecting and evaluating them] Drevnie redkometal'no-titanovye rossypi, metody ikh poiskov i otsenki. Moskva, Nedra, 1964. 169 p. (MIRA 17:12)

KAZARINOV, N.A. [Kazarinov, M.O.]; DZYUBA, N.P.

New method for quantitative determination of khellin in
anhydrous solvents. Farmatsev. zhur. 18 no.4:39-43 '63.

(MIRA 17:7)

1. Khar'kovskiy nauchno-issledovatel'skiy khimiko-farmatsevticheskiy
institut.

L 1795-66

ACCESSION NR: AP5017529

UR/0243/85/000/007/0057/0060

815.711.5-014.3:543.24

AUTHOR: Kazarinov, N. A.; Dzyuba, N. P.

TITLE: Determination of pharmaceutical preparations containing a carbonyl group by titration with nonaqueous solvents. Report 3. Analysis of cardiac glycosides

AOOURCE: Meditsinskaya promyshlennost' SSSR, no. 7, 1965, 57-60

TOPIC TAGS: circulatory drug, quantitative analysis, chemical identification, analytic chemistry

ABSTRACT: Earlier controls of these preparations were conducted with inconvenient biological methods. A quantitative chemical method described in earlier publications was applied to glycosides containing an aldehyde group at C₁₀, in particular to cymarin, erysimin, semisynthetic and natural convallatoxin and Corelborin II (Helleborus glycoside). It consists in reacting the carbonyl group with hydroxylamine HCl, forming the corresponding oxime and binding the liberated HCl with diethylamine whose excess is titrated off with perchloric acid in me-

Card 1/2

L 1795-66

ACCESSION NR: AP5017529

4
thanol in the presence of the thymol blue indicator. Chromatographic tests showed that the results agreed with data from biological analysis, but the results obtained by chemical analysis were more reproducible and more exact. The chemical analysis method is therefore recommended for both laboratory work and analysis of standards. The semisynthetic convallatoxin was prepared by V. T. Chernobay. Orig. art. has: 7 formulas and 1 table

55
ASSOCIATION: Khar'kovskiy nauchno-issledovatel'skiy khimiko-farmatsevticheskiy institut (Kharkov Chemical-Pharmaceutical Scientific Research Institute)

SUBMITTED: 25Apr64

ENCL: 00

SUB CODE: LS, G-C

NR REF SOV: 013

OTHER: 004

mlb
Card 2/2

KAZARINOV, N.A. [Kazarynov, M.O.]; DZYUBA, N.P.

Relation of the rate of the formation of oximes of cardiac glycosides to their structure. Farmatsev. zhur. 20 no.5:28-31 '65.
(MIRA 18:11)

1. Khar'kovskiy nauchno-issledovatel'skiy khimiko-farmatsevticheskiy institut. Submitted January 26, 1965.

KAZARINOV, N.D.

Scalar diffraction problem for an elliptic cylinder and an ellipsoid of revolution. Dokl.AN SSSR 138 no.4:759-762 Je '61.

(MIRA 14:5)

1. Matematicheskiy institut imeni V.A.Steklova AN SSSR. Predstavleno akademikom A.A.Dorodnitsynym.

(Potential, Theory of)

(Boundary value problem)

BING, R.G.; KAZARINOV, N.D. (Madison, Wisconsin, SSHA); KAZHDAN, I.A.,
(studentka 4-go kursa); MAS'KO, S.S. (studentka 4-go kursa); DORFMAN,
A.G. (Gor'kiy); KUZHEL', A.V. (Uman'); SKOPETS, Z.A. (Yaroslavl');
TELESIN, Yu.Z. (Moskva)

Brief notes. Mat. pros. no. 6: 205-216 '61.

(MIRA 15:3)

1. Moskovskiy gosudarstvennyy pedagogicheskiy institut imeni Lenina
(for Kazhdan, Mas'ko).

(Mathematics—Problems, exercises, etc.)

33238

S/089/62/012/002/010/013

B102/B138

26.2264
21.6000

AUTHORS: Kazarinov, N. M., Matveyev, O. A., Ryvkin, S. M., Solov'yev, S. M., Strokan, N. B., Tarkhin, D. V.

TITLE: Investigation of semiconductor spectrometer counters for measuring fragment energies

PERIODICAL: Atomnaya energiya, v. 12, no. 2, 1962, 153 - 154

TEXT: U^{235} fission fragment energy was measured by semiconductor counters developed at the fiziko-tekhnicheskiy institut im. A. F. Ioffe (Physicotechnical Institute imeni A. F. Ioffe). The surface-barrier junction of these counters was produced by spraying gold onto an n-type silicon plate. These counters, which were studied earlier by the authors (Atomnaya energiya, 11, no. 3, 217, 1961), were found to be well suited for alpha spectrometry (resolution 0.5% for $E_{\alpha} = 5.5$ Mev). The volume charge region was about 60μ for maximum voltage, much greater than the fragment range in silicon. Fragment energy was measured with a 0.5 mm Al target, placed in a thin-walled aluminum vacuum chamber. The target had a vacuum-sprayed layer of UF_4 , enriched in U^{235} to 92.8%. Diameter of the Card 1/3

53238

S/089/62/012/002/010/013

B102/B138

Investigation of semiconductor v..

layer was 1.2 cm, and the total weight was 120 μ g. The silicon counter was placed 1.5 cm below the target to avoid being hit by the neutron beam collimated into the chamber. The counter pulses were fed to a preamplifier and thence to a 100-channel analyzer. The fragment energy spectra thus measured differed considerably from those obtained from time-of-flight measurements. This was found to be due to energy losses in the counter surface, which were strongly dependent on the angle of incidence of the fragments. As the fragments lose most of their energy in the first part of their path this effect was much higher for them than for alphas ✓

Special counters of 16 mm² area were produced with a thinner layer of gold and the energy spectrum was measured again and compared as before. This time the shape was the same, with a difference of about 7 Mev in absolute values. This is attributed partly to energy losses in the fissile layer, and partly to the energy being carried away by fission neutrons. In the Au layer losses do not exceed 1 Mev. Apart from other advantages the silicon counters yield better results than e. g. ionization chambers. There are 2 figures and 5 references: 1 Soviet and 4 non-Soviet. The four references to English-language publications read as follows: W. Stein, Card 2/3

37103

8/056/62/042/004/015/037
B152/B102

21.1000

AUTHORS: Blinov M. V., Kazarinov N. M., Protopopov A. N.

TITLE: Study of the energy and angular distributions of neutrons emitted in thermal-neutron induced U^{235} fission

PERIODICAL: Zhurnal eksperimental'noy i teoreticheskoy fiziki, v. 42, no. 4, 1962, 1017-1021

TEXT: The authors measured the energy distribution of prompt neutrons emitted by thermal neutrons in U^{235} fission for the angles 0° , 45° , and 90° to the flight direction of fission fragments. The fragments were recorded by a xenon-filled ($p = 1.5$ atm) scintillation counter. An aluminum foil with a thin U^{235} layer (~ 2 mg/cm²) was attached to this counter. The most probable angle of departure of the fragments was determined by collimators on these layers. Stilbene scintillation detectors for detecting the fission neutrons were placed at a certain distance from the uranium layer at various angles to the flight direction of the fragments. The neutron energy was calculated from the time of flight between the two counters. The time of flight was determined by a

Card 1/3

X

Study of the energy and angular ...

S/056/62/042/004/015/037
B152/B102

100-channel time analyzer. The coincidences were taken with $\Phi 33$ -33 (FEU-33) photomultipliers. The half width of the coincidence distribution was $6 \cdot 10^{-10}$ sec for Co^{60} gamma quanta for two pairs of these multipliers. The neutron energy threshold still recorded was about 100 kev. The most important part of this time scale was calibrated on the basis of the flight time of gamma quanta for different paths. The remaining part of the scale was gauged with calibrated pieces of the PK-2 (RK-2) cable. The time resolution in the experiments was $5 \cdot 10^{-9}$ sec. The measurements gave the following ratios as relative neutron emission probability: $N(0^\circ):N(45^\circ):N(90^\circ) = (5.7 \pm 0.2):(2.9 \pm 0.1):1$. The energies found in this study are harder for 45° and 90° , and much harder for 0° than those found by V. N. Nefedov (ZhETF, 38, 1657, 1960). The values of the present paper do not agree with the calculations of Yu. A. Vasil'yev et al. (Atomn. energ. 2, 449, 1960). The same was also observed by Vasil'yev for the 14-Mev neutron induced fission of uranium (Yu. A. Vasil'yev et al., ZhETF, 38, 671, 1960). The studies were made with the reactor of the Physicotechnical Institute of the Academy of Sciences USSR. S. M. Solov'yev is thanked for special measurements, A. D. Kolchin, L. I. Radayev, V. V. Pikunov and A. G. Roshchin for technical aid. There are 3 figures.

Card 2/3

43364

S/056/62/043/005/012/058
B102/B104

24.6410

AUTHORS: Blinov, M. V., Kazarinov, N. M., Protopopov, A. N.,
Shiryayev, B. M.

TITLE: The angular anisotropy of γ -quanta accompanying the U^{235}
fission

PERIODICAL: Zhurnal eksperimental'noy i teoreticheskoy fiziki, v. 43,
no. 5(11), 1962, 1644-1648

TEXT: The angular anisotropy of γ -quanta emitted in thermal-neutron induced fission of U^{235} was measured under optimum geometry. The γ -quantum detector, a stilbene crystal and an $\Phi 34-33$ (FEU-33) photo-multiplier, was lead-shielded (10 cm) in order to reduce the background effect. The effects of prompt neutrons were eliminated by a multi-channel time analyzer (resolution $3-5 \cdot 10^{-9}$ sec). In previous measurements made with a gas scintillation counter and a copper collimator the anisotropy was found to be 25-30%. Control measurements showed that this high value can be somewhat reduced if account is taken of the γ -quantum
Card 1/3

The angular anisotropy of ...

S/056/62/043/005/012/058
B102/B104

absorption both of the collimator and of the Al backing of the target. In the following experiments these effects were eliminated with the help of a special arrangement (Fig. 1). The fragments were detected by the flashes in a thin scintillation foil. The time distributions of the gamma counts taken at 0 and 90° angles to the chamber axis show a peak with a half-width of $8 \cdot 10^{-9}$ sec. Records gave $19.5 \cdot 10^6$ fission events (13,063 quanta) under 0° and $19.8 \cdot 10^6$ (12,069 quanta) under 90° to the axis. Thus after all corrections the anisotropy amounts to $\frac{W(0)-W(90)}{W(90)} = (12 \pm 2)\%$, i. e. the γ -emission in the direction of flight of fragments is higher by (12±2)% than perpendicular thereto. Assuming it is the fragment's angular momentum that causes the anisotropy its value can be estimated. For $L \gg 2$, $j \sim 25 - 30$, for $L=3$, $j \sim 15 - 20$. The measurements described have been made at the reactor of the FTI AN SSSR. There are 2 figures.


SUBMITTED: June 18, 1962

Card 2/3

The angular anisotropy of ...

8/056/62/043/005/012/058
B102/B104

Fig. 1. Apparatus for measuring the γ -ray intensity along the chamber axis. Legend: 1 - neutron beam; 2 - FEU-33, 3 - Terphenyl film, 4-uranium oxide layer (97.9% U^{235}), 5 - vacuum chamber, 6 - stilbene crystal, 7 - lead shield.



Card 3/3

ACCESSION NR: AP4018361

S/0120/64/000/001/0040/0045

AUTHOR: Blinov, M. V.; Kazarinov, N. M.

TITLE: Fission-neutron spectrometer

SOURCE: Pribery* i tekhnika eksperimenta, no. 1, 1964, 40-45

TOPIC TAGS: neutron, fission neutron, spectrometer, fission neutron spectrometer, gas scintillation counter, ionization chamber, neutron spectroscopy

ABSTRACT: A spectrometer for measuring fission neutrons within 200 kev to 7 Mev by the transit-time method is described. Time intervals were measured by converting them into a pulse-height distribution which was recorded by an AI-100-1 multichannel pulse-height analyzer. The time resolution of the instrument, determined from the half-width of γ - γ coincidence of Co^{60} , for 100% efficient recording of γ quanta is 2 nanosec; the resolution for a 10% efficiency is

Card 1/2

ACCESSION NR: AP4018361

0.7 nanosec. The apparatus resolution of the time analyzer is under 10^{-10} sec. A multilayer gas scintillation counter, a semiconductor detector, and an ionization chamber were used to detect fragments. Half-widths of the time distribution of fission gamma-rays obtained with the above detectors were 3, 3-4, and 10 nanosec, respectively. The spectrometer proved to be a highly stable instrument: a time scale drift of only 3 nanosec was noticed after a two-month period of daily operation. "The authors are indebted to A. N. Protopopov for discussing the results of this work, and to A. G. Roshchin for assembling the equipment." Orig. art. has: 9 figures.

ASSOCIATION: none

SUBMITTED: 14Mar63

DATE ACQ: 18Mar64

ENCL: 00

SUB CODE: NS

NO REF SOV: 008

OTHER: 003

Card 2/2

L 27896-68 INT(ENR/ENR/1) TEMP(1) INT(1) CI

ACCESSION NO: AP4012268

5/0080/64 014 000000

AUTHOR: Kharin, V. I. A. Kharin, V. I.

TITLE: Surface-barrier silicon counters in a neutron flux of
frequency

SOURCE: Atomnaya energiya, v. 16, no. 1, 1968, p. 100.

TOPIC TAGS: silicon counter; counter irradiation; surface-barrier counter

ABSTRACT: The authors investigated the changes in some characteristics of

TOP SECRET

SECRET

1.10544

ALICE, 1.10544

pulses corresponding to the y-quanta were supplied from a
photomultiplier to the time converter through an attenuator

App. 1.10544

SUBMITTED: 21 Nov 65

ATT. PRESS: 31.1

END




Fig. 1 Block diagram of the fast-slow coincidence-circuit system

D and D₁ - Amplitude discriminator;
 1 - coincidence circuit; 2 - fast
 time amplitude analyzer; 3 - slow
 time amplitude analyzer; 4 - fast
 time amplitude analyzer; 5 - slow
 time amplitude analyzer; 6 - coincidence circuit

Card 3

ACCESSION NR: AP4025950,

S/0056/64/046/003/1139/1141

AUTHORS: Elinov, M. V.; Kazarinov, N. M.; Protopopov, A. N.

TITLE: Angular and energy characteristics of emission of U-235 fission neutrons

SOURCE: Zhurnal eksperimental'noy i teoreticheskoy fiziki, v. 46, no. 3, 1964, 1139-1141

TOPIC TAGS: uranium 235, uranium fission, fission neutron, emission spectrum, energy distribution, angular distribution, cascade evaporation theory

ABSTRACT: In contrast to earlier work by the authors (ZhETF v. 42, 1017, 1962) and by V. N. Nefedov (ZhETF v. 38, 1657, 1960), the present study is devoted to an experimental determination of the c.m.s. energy spectrum of the neutrons from thermal neutron fission of U^{235} . This emission spectrum is then used to calculate the energy and angu-

Card 1/2

ACCESSION NR: AP4025950

lar distributions in the laboratory system. The results of these calculations are compared with experimental distributions, which are measured in more detail than before. Certain discrepancies between the experimental and calculated data are discussed, but in spite of the discrepancies it is concluded that the overwhelming majority of the neutrons (~90%) emitted, following thermal-neutron fission of U^{235} are emitted in an ordinary cascade evaporation process from fully accelerated fragment nuclei. The conclusions are drawn from the fact that the experimental neutron-emission spectrum agrees with calculations based on the neutron cascade evaporation theory (K. J. LeCouteur and D. W. Lang, Nuclear Physics, v. 13, 32, 1959). A detailed report will be published in "Atomnaya energiya." Orig. art. has: 1 figure.

ASSOCIATION: None

SUBMITTED: 13Aug63

DATE ACQ: 16Apr64

ENCL: 00

SUB CODE: PH, NS
Card 2/2

NR REF SOV: 002

OTHER: 001

Card 1 of 2

ASSOCIATION. NOTE

Card 2/3

24.7700 (1035, 1043, 1469)

S/056/61/040/003/025/031
B108/B209

AUTHORS: Kazarinov, R. F., Konstantinov, O. V.

TITLE: Dispersion theory of high-frequency exciton conductivity
in a crystal

PERIODICAL: Zhurnal eksperimental'noy i teoreticheskoy fiziki, v. 40,
no. 3, 1961, 936-942

TEXT: The authors employ the graph technique suggested in Ref. 5
(O. V. Konstantinov, V. I. Perel'. ZhETF, 32, 197, 1960) for the
calculation of high-frequency conductivity. They discuss direct transitions
(without phonons) in which the energy maximum of the valency band and the
minimum of the conduction band do not coincide in the momentum space.
 K_1 are the threshold points for direct transition at the cutoff frequency
 ω_0 . The excitons forming at these points on transition are equivalent,
and their spectra coincide when the wave vector of light is neglected.
These excitons are also formed when electrons of the mean velocity

Card 1/28

1115

Dispersion theory of ...

S/056/61/040/003/025/031
B108/B209

$\vec{v}_1 = \vec{v}_p / \vec{p}$ $\vec{p} = \vec{k}_1$ absorb light with the Doppler frequency shift

$\Delta \omega_1 = \vec{v}_1 \cdot \vec{k}_1$, where \vec{k}_1 denotes the wave vector of light. The problem of high-frequency conductivity is solved for the simple model of an electron gas in an alternating field that secures electron-electron interaction at energies that are much less than the width of the forbidden band. The Hamiltonian H of such a system is given by

$$H = H_0 + U; \quad H_0 = \sum_{j\vec{p}} \epsilon_{j\vec{p}} a_{j\vec{p}}^+ a_{j\vec{p}}, \quad U = \sum_{j_1\vec{p}_1, j_2\vec{p}_2, j_3\vec{p}_3, j_4\vec{p}_4} U_{j_1\vec{p}_1, j_2\vec{p}_2, j_3\vec{p}_3, j_4\vec{p}_4} a_{j_1\vec{p}_1}^+ a_{j_2\vec{p}_2}^+ a_{j_3\vec{p}_3} a_{j_4\vec{p}_4}, \quad (1)$$

$$U_{j_1\vec{p}_1, j_2\vec{p}_2, j_3\vec{p}_3, j_4\vec{p}_4} = \sum_{\gamma, \gamma', m} \frac{4\pi e^2}{\epsilon \gamma^2 V} \delta_{\gamma+\vec{p}_1, \vec{p}_2+\vec{p}_3+\vec{p}_4} \delta_{\gamma'+\vec{p}_1, \vec{p}_2+\vec{p}_3+\vec{p}_4} \Gamma_{j_1\vec{p}_1}(\vec{p}_1, \vec{p}_2, \gamma) \Gamma_{j_2\vec{p}_2}(\vec{p}_2, \vec{p}_3, -\gamma),$$

$$\Gamma_{j\vec{p}}(\vec{p}_1, \vec{p}_2, \gamma) = \int_{\vec{v}} e^{i\vec{v}\cdot\vec{x}} \psi_{j\vec{p}}^*(\vec{x}) \psi_{j\vec{p}}(\vec{x}) d\vec{x}.$$

where $a_{j\vec{p}}^+$ and $a_{j\vec{p}}$ are the production and annihilation operators of an electron in a state with the band j and the quasi-momentum \vec{p} , which are described by the Bloch wave function $\psi_{j\vec{p}}(\vec{x})$; $\epsilon_{j\vec{p}}$ denotes the energy of

Card 2/28

22145

S/056/61/040/G03/025/031
B108/B209

Dispersion theory of ...

this state, V the normalized volume, V_0 the volume of a lattice cell, and \vec{b}_m and \vec{b}_n are multiples of the vector of the reciprocal lattice; the dielectric constant ϵ is assumed to be a scalar. With formula (4a) from Ref. 5, the authors found the correction to the density matrix $\rho_{j\vec{p},j'\vec{p}'}$ to be proportional to the strength of the electric field applied $E_\mu(\vec{x},t) = E_\mu(\vec{x},s)\exp(i\vec{x}\vec{\omega} + st)$, where $s = -i\omega + \nu$; $\vec{\omega}$ and ω denoting wave vector and frequency of light, ν an adiabatic parameter:

$$\rho_{j\vec{p},j'\vec{p}'}(t) = E_\mu(x,s) e^{i\mu} \sum_{\vec{k},\vec{k}'} G_{\vec{k},\vec{k}'}^{j\vec{p},j'\vec{p}'}(s,\beta) \int e^{i\vec{k}\vec{x}} j_\mu(x)_{\vec{k},\vec{k}'} dx; \quad (2a) \quad (2a)$$

$$G_{\vec{k},\vec{k}'}^{j\vec{p},j'\vec{p}'}(s,\beta) = Z^{-1} \int_{-\infty}^0 e^{s\tau} d\tau \int_0^\beta d\lambda \text{Sp} \left\{ e^{-H\tau} \exp \left[\frac{H(\tau + i\hbar\lambda)}{i\hbar} \right] a_{j\vec{p}}^\dagger a_{j'\vec{p}'} \exp \left[-\frac{H(\tau + i\hbar\lambda)}{i\hbar} \right] a_{\vec{k}}^\dagger a_{\vec{k}'} \right\},$$

$$(j_\mu(x))_{\vec{k},\vec{k}'} = (e/2m) [\psi_{\vec{k}}^*(x) \hat{p}_\mu \psi_{\vec{k}'}(x) - \psi_{\vec{k}'}(x) \hat{p}_\mu \psi_{\vec{k}}^*(x)], \quad (2b) \quad (2c)$$

$$(2a) \quad (2b) \quad (2c).$$

Card 3/9 8

Dispersion theory of ...

22145
S/056/61/040/003/025/031
B108/B209

X

$Z = \sum_{\mathbf{k}} e^{-\beta \epsilon_{\mathbf{k}}}$, $\beta = T^{-1}$, T is the absolute temperature in energy units. The quantities $G_{\mathbf{k}, \mathbf{j}\mathbf{p}}^{i' \mathbf{k}', j' \mathbf{p}'}(s, \beta)$ are determined by the graph method of Ref. 5.

The case under consideration is characterized by one-electron states which are represented by Bloch wave functions. A combination of indices j and \mathbf{p} corresponds to every line. Fig. 1 shows a peak illustrating electron-electron interaction corresponding to the factor subsequent to Fig. 1. The quasi-momentum of each line does not exceed half the vector of the reciprocal lattice by its absolute amount. Therefore, the quantity $(\mathbf{b}_n - \mathbf{b}_m)$ does not exceed the vector of the reciprocal lattice. The lines correspond to the factors $1 - n_{j\mathbf{p}}$ or $n_{j\mathbf{p}}$, where $n_{j\mathbf{p}}$ is the Fermi function.

The authors examine the case in which $T \ll \epsilon_0$; $n_{j\mathbf{p}}$ equals unity when j is the index of a completely filled band, and equals zero when j refers to a band that is incompletely filled. The authors then discuss the various perturbation-theoretical approximations. Fig. 2 shows the case of

Card 4/28

Dispersion theory of ...

S/056/61/040/003/025/031
B108/B209

zeroth approximation. Since the first-approximation terms have a common factor (Fig. 4b), one may introduce the quantity $F_{\vec{k}\vec{p}}^{\vec{k}'\vec{p}'}(s)$, where

$$G_{\vec{k}\vec{p}}^{\vec{k}'\vec{p}'}(s, \sigma) = \frac{1}{\sigma} \frac{\delta_{\vec{k}+\vec{p}, \vec{k}'+\vec{p}'}}{\sigma + \epsilon_{\vec{k}} - \epsilon_{\vec{p}'}} F_{\vec{k}\vec{p}}^{\vec{k}'\vec{p}'}(s). \quad (5)$$

and

$$F_{\vec{k}\vec{p}}^{\vec{k}'\vec{p}'}(s) = \delta_{\vec{k}\vec{p}} [s + i\hbar^{-1}(\epsilon_{\vec{p}'} - \epsilon_{\vec{p}})]^{-1} - \\ - (i\hbar)^{-1} \sum_{\vec{\gamma}} F_{\vec{k}, \vec{p}+\vec{\gamma}}^{\vec{k}', \vec{p}'+\vec{\gamma}}(s) \frac{4\pi e^2 \epsilon^{-1}}{V\gamma^4} \Gamma_{\alpha}(\vec{p} + \vec{\gamma}, \vec{p}, \vec{\gamma}) \times \\ \times \Gamma_{\alpha}(\vec{p}', \vec{p}'+\vec{\gamma}, -\vec{\gamma}) [s + \frac{i}{\hbar}(\epsilon_{\vec{p}'} - \epsilon_{\vec{p}})]^{-1}. \quad (6)$$

With the identity $R(\vec{p}, \vec{k}') \equiv F_{\vec{k}', \vec{p}+\vec{z}}^{\vec{k}', \vec{p}'+\vec{z}}(s)$, the authors obtain the following final result:

Card 5/28

22145

Dispersion theory of ...

S/056/61/040/003/025/031
B108/B209

X

$$R(p, k') = \begin{cases} \frac{1}{V} \sum_n \frac{\varphi_n(p - K_1 - \lambda) \varphi_n^*(k' - K_1 - \lambda)}{s + i(\omega_n + v'z)}, & p, k' \sim K_1 + \lambda, \\ \frac{1}{V} \sum_n \frac{\varphi_n(p - K_f - \lambda) \varphi_n^*(k' - K_f - \lambda)}{s + i(\omega_n + v'z)}, & p, k' \sim K_f + \lambda. \end{cases} \quad (13) \quad (13),$$

where $\hbar\omega_n = E_0 + \frac{1}{2} M_{\alpha\beta}^{-1} \vec{r}_{\alpha\beta}^2 + \epsilon_n$; the wave function $\varphi_n(\vec{q})$ of an exciton in the n -th state satisfies the equation

$$\mu_{\alpha\beta}^{-1} \frac{\hbar^2}{2} q_{\alpha} q_{\beta} \varphi_n(q) - \sum_{\gamma} \frac{4\pi e^2 s^{-1}}{\gamma^3} \frac{1}{V} \varphi_n(q + \gamma) = \epsilon_n \varphi_n(q) \quad (14) \quad (14).$$

With Fourier transformation, this leads to the Schrödinger equation for the motion of the exciton. With Eqs. (13), (8), (5), and (2a), the authors obtain the following expression for high-frequency conductivity:

Card 6/98

21145

Dispersion theory of ...

S/056/61/040/003/025/031
B108/B209

$$\sigma_{\mu\nu}(\mathbf{x}, \omega) = E_0^{-1} (2\pi)^{-3} \sum_{n,l} \int d\mathbf{p} d\mathbf{k}' \frac{\varphi_n(\mathbf{p} - \mathbf{K}_l - \lambda) \varphi_n^*(\mathbf{k}' - \mathbf{K}_l - \lambda)}{s + i(\omega_n + v^2 \mathbf{x})} j_\mu(\mathbf{k}', \mathbf{x}) j_\nu^*(\mathbf{p}, \mathbf{x}).$$

(16) (16).

For allowed transitions where $j_\mu(\vec{K}_1, 0) \neq 0$, this expression is simplified because for long-range excitons, $\varphi(\vec{p} - \vec{K}_1 - \vec{\lambda})$ is non-vanishing only in the immediate neighborhood of the point $\vec{K}_1 + \vec{\lambda}$ in the \vec{p} space. Neglecting the dependence of $j_\mu(\vec{k}', \vec{\lambda})$ on \vec{k}' , one obtains

$$\chi_{\mu\nu}(\mathbf{x}, \omega) = \frac{e^2}{\hbar} \sum_n |\tilde{\varphi}_n(0)|^2 \sum_l \frac{r_{\nu}^*(K_l) r_{\mu}^*(K_l)}{(\omega - \omega_n - v^2 \mathbf{x}) + i\nu}, \quad (17)$$

$$r_{\nu}^*(K_l) = \int \psi_{\nu K_l}^*(r) r^\nu \psi_{\nu K_l}(r) dr. \quad (17)$$

for high-frequency exciton polarizability. This formula is analogous to

Card 7/9 8

Dispersion theory of ...

22145
S/056/61/040/003/025/031
B108/B209

that for a gas. The authors thank L. E. Gurevich and V. I. Perel' for numerous discussions. There are 5 figures and 7 references: 3 Soviet-bloc and 4 non-Soviet-bloc. The two references to English-language publications read as follows: Ref. 1: R. I. Elliott. Phys. Rev., 108, 1383, 1957; Ref. 7: J. J. Hopfield. Phys. Rev., 112, 1555, 1958.

ASSOCIATION: Leningradskiy fiziko-tekhnicheskii institut Akademii nauk SSSR (Leningrad Institute of Physics and Technology of the Academy of Sciences USSR)

SUBMITTED: October 19, 1960



Рис. 1
Fig. 1

$$\pm (in)^{-1} \sum_{mn} \frac{4\pi e^2 b^{-1}}{V(p_2 - p_1 + b_n)^2} \Gamma_{1/2}(p_1, p_2, p_2 - p_1 + b_n) \times \\ \times \Gamma_{1/2}(p_2, p_1, p_2 - p_1 - b_m) \delta_{p_1 + p_2 - p_1 - b_m - b_n}.$$

Card 8/98

26419
S/056/61/041/001/014/021
B102/B214

24.7700

AUTHORS: Ipatova, I. P., Kazarinov, R. F.

TITLE: Faraday effect on excitons

PERIODICAL: Zhurnal eksperimental'noy i teoreticheskoy fiziki, v. 41,
no. 1(7), 1961, 209-210

TEXT: This paper gives a theoretical investigation of the rotation of the plane of polarization in the neighborhood of exciton absorption lines; the angle of this rotation depends on the effective mass and the radius of the exciton. In the neighborhood of the exciton absorption lines (corresponding to the transition into the p state) the rotation of the plane of polarization (Faraday effect) may be expected to be large. In cubic crystals the angle of rotation is expressed by the component of the boundary vector G lying in the H direction ($H \parallel z$): $\varphi = (\pi d / \lambda) G_z / \epsilon$, where d is the thickness of the sample, ϵ the dielectric constant without magnetic field, and λ the wavelength of light. G is determined by the asymmetric part of the dielectric constant: $G_\gamma = \delta_{\gamma\mu\nu} \epsilon_{\mu\nu}$. Therefore,

Card 1/4

26419

S/056/61/041/001/014/021

B102/B214.

Faraday effect on excitons

the problem consists in the calculation of $\epsilon_{\mu\nu}(\vec{H})$. For Mott's excitons only transitions to the s-state are permitted, those to the p-state are forbidden. Since, however, the s-state transitions show no Faraday effect, the less intensive forbidden transitions are considered. For the exciton conductivity one has:

$$\sigma_{\mu\nu} = \sum_{n;m=0,\pm 1} T_{\mu\nu}^{nm} / [\gamma - i(\omega - \omega_n - \Omega m)], \quad (3),$$

where ω is the frequency of light, $\Omega = eH/2Mc$, M the reduced mass of the exciton, ω_n the hydrogen-like energy level, γ the width of the exciton line, m the magnetic quantum number, $T_{\nu\mu}^{nm}$ a tensor whose real part is related with the oscillator force of the transition and the imaginary part with the rotation of the plane of polarization:

$$T_{\mu\nu}^{nm} = \frac{i}{E_0} \sum_l \left(\frac{\partial J_{\nu}^{(l)}}{\partial k_{\alpha}} \right)_{K_l} \left(\frac{\partial J_{\mu}^{(l)}}{\partial k_{\beta}} \right)_{K_l} \left(\frac{\partial \Psi_{n1m}}{\partial x_{\beta}} \right)_{x=0} \left(\frac{\partial \Psi_{n1m}}{\partial x_{\alpha}} \right)_{x=0}. \quad (4)$$

E_0 is the minimum frequency of the transition to the ground state, K_l is the point in the momentum space corresponding to this transition, Ψ_{n1m}

Card 2/4

Faraday effect on excitons

26419
S/056/61/041/001/014/021
B102/B214

the hydrogen-like wave function of the p-state of the exciton, and

$$J_{\nu}^{\mu}(k) = J_{\nu}^{\mu}(k) = e \int d^3r u_{\nu}(r) \hat{v}_{\nu} u_{\mu}(r)$$

the matrix element of the band-to-band current calculated from Bloch's wave function. With $T_{\mu\nu}^{nm} = T_{\nu\mu}^{n-m}$; $T_{\mu\nu}^{nm} = (T_{\nu\mu}^{nm})^*$ and the assumption that $|\omega_n - \omega| \ll \gamma$, one obtains

$$-i\epsilon_{\mu\nu}^{(n)} = \frac{2\pi}{\omega} (\sigma_{\mu\nu} - \sigma_{\nu\mu}) = -i \frac{4\pi}{\omega} \sum_n \text{Im}(T_{\mu\nu}^{nn}) \frac{\Omega}{\gamma^2 + (\omega - \omega_n)^2} \quad (6)$$

The angle of rotation near the line with $n = 2$ is given by:

$$\varphi = \frac{\pi}{2} \frac{1}{\epsilon} \left(\frac{d}{\lambda} \right) \left(\frac{a}{2r_0} \right)^2 \frac{\Omega\omega}{\gamma^2 + (\omega - \omega_n)^2} \beta_{xy} \quad (7)$$

where $\beta_{xy} \sim 1$. A rough numerical estimate of φ for the yellow exciton series in Cu_2O crystal ($\epsilon=10$) gives: $\varphi \sim 0.5^\circ$ for $H = 10^3$ gauss,

$|\omega - \omega_n| \sim \gamma \sim 10$, $r_0 \leq 30a$, and $d = 500\mu$. From the angle of rotation

Card 3/4

Faraday effect on excitons

26419
S/056/61/041/001/014/021
B102/B214

which is always measurable, one can obtain the radius of the exciton if its reduced mass is known. The authors thank L. E. Gurevich, O. V. Konstantinov, and G. M. Eliashberg for discussions. There are 5 references: 4 Soviet-bloc and 1 non-Soviet-bloc.

ASSOCIATION: Leningradskiy fiziko-tekhnicheskii institut Akademii nauk SSSR (Leningrad Institute of Physics and Technology of the Academy of Sciences, USSR)

SUBMITTED: January 31, 1961

card 4/4

24.7100

35106

S/056/62/042/002/021/037
B108/B102

AUTHORS: Kazarinov, R. F., Skobov, V. G.

TITLE: Theory of nonlinear galvanomagnetic phenomena
in semiconductors

PERIODICAL: Zhurnal eksperimental'noy i teoreticheskoy fiziki,
v. 42, no. 2, 1962, 1047 - 1053

TEXT: The current through a semiconductor in crossed magnetic and electrical fields is studied under both classical and quantum limit conditions. The energy imparted to the electrons on scattering is shown to be greater than they can impart to the lattice. This causes the temperature of the electron gas to rise. In the quantum case, the electron distribution on scattering is Boltzmannian; the effective electron temperature is proportional to the square of the electrical field strength. The transverse current density is proportional to E^2/H^3 in the classical case, and proportional to H^3/E^2 in the quantum limit. This result was obtained

Card 1/2

Theory of nonlinear galvanomagnetic ...

S/056/62/042/002/021/037
B108/B102

on the assumption that the conduction electron concentration was constant. The dependence of resistivity on the current density j is similar for scattering of electrons from phonons and neutral impurities and from ionized impurities. In the quantum limit resistivity decreases as the cube j . L. E. Gurevich is thanked for having suggested the theme, V. I. Perel' and G. M. Eliashberg for discussion of the work. There are 1 figure and 3 references: 1 Soviet and 2 non-Soviet. The English-language reference reads as follows: E. N. Adams, T. D. Holstein, J. Phys. Chem. Solids, 10, 254, 1959.

ASSOCIATION: Leningradskiy fiziko-tekhnicheskii institut Akademii nauk SSSR (Leningrad Physicotechnical Institute of the Academy of Sciences USSR)

SUBMITTED: September 5, 1961

Card 2/2

24.1800 6.8000

3556,
S/056/62/042/003/045/049
B108/B102

AUTHORS: Kazarinov, R. F., Skobov, V. G.

TITLE: Possibility of amplifying ultrasound in semimetals in an electric field

PERIODICAL: Zhurnal eksperimental'noy i teoreticheskoy fiziki, v. 42, no. 3, 1962, 910-912

TEXT: The amplification of ultrasound by conduction electrons in an electric field is discussed. Amplification is explained as follows: When no electric field is present, the electrons will absorb acoustic energy Q_0 . This leads to an electron-acoustic current \vec{J} proportional to Q_0 . The sonic energy absorbed by the electrons per unit time is therefore $Q = Q_0 + \vec{J}\vec{E}$ (linear approximation with respect to E). The sound will be amplified by the electrons when \vec{E} is such that $Q < 0$. When one sound quantum is absorbed, the electron velocity will change by the amount $\hbar\kappa/m$ (κ - wave vector of sound, m - effective mass of electron). During the time τ between two collisions the electron is shifted by $\Delta = \hbar\kappa\tau/m$. In

Card 1/2